

Mammary-specific deletion of parathyroid hormone-related protein preserves bone mass during lactation

Joshua N. VanHouten, ... , Andrew C. Karaplis, John J. Wysolmerski

J Clin Invest. 2004;113(3):492-492. <https://doi.org/10.1172/JCI19504C1>.

Corrigendum

Endocrinology

Original citation: *J. Clin. Invest.* 112:1429–1436(2003). doi:10.1172/JCI19504. Citation for this erratum: *J. Clin. Invest.* 112:492 (2004). doi:10.1172/JCI19504C1. A number of errors were included in Table 1. The correct version is as follows: Table 1 Histomorphometric analysis of tibias from lactating BLG-Cre/PTHrPlox^{-/-} mice and PTHrPlox^{-/-} controls

Find the latest version:

<https://jci.me/19504C1/pdf>



Mammary-specific deletion of parathyroid hormone–related protein preserves bone mass during lactation

Joshua N. VanHouten, Pamela Dann, Andrew F. Stewart, Christine J. Watson, Michael Pollak, Andrew C. Karaplis, and John J. Wysolmerski

Original citation: *J. Clin. Invest.* **112**:1429–1436 (2003). doi:10.1172/JCI200319504.

Citation for this corrigendum: *J. Clin. Invest.* **113**:492 (2004). doi:10.1172/JCI200419504C1.

A number of errors were included in Table 1. The correct version is as follows:

Table 1

Histomorphometric analysis of tibias from lactating *BLG-Cre/PTHrP^{lox/-}* mice and *PTHrP^{lox/-}* controls

	OcS/BS (%)	NOc/BPm (no./mm)	ObS/BS (%)	NOb/BPm (no./mm)	BFR/TV (%/yr)	BV/TV (%)	TbTh (μ m)	TbSp (μ m)	TbN (no./mm)
<i>PTHrP^{lox/-}</i>	7.0 \pm 1.0	2.3 \pm 0.3	18.2 \pm 2.3	21.8 \pm 3.4	372.7 \pm 36.2	6.1 \pm 0.5	25.3 \pm 2.1	402.2 \pm 48.4	2.46 \pm 0.27
<i>BLG-Cre/PTHrP^{lox/-}</i>	3.0 \pm 0.5 ^A	1.0 \pm 0.1 ^A	18.7 \pm 1.9	26.0 \pm 3.0	219.5 \pm 30.0 ^B	6.8 \pm 1.2	21.9 \pm 2.1	379.9 \pm 96.1	3.28 \pm 0.62

^A*P* < 0.005; ^B*P* < 0.05. Static and dynamic histomorphometric analysis of tibias from day 12 lactating *BLG-Cre/PTHrP^{lox/-}* mice (*n* = 7, static; *n* = 4, dynamic) and *PTHrP^{lox/-}* littermate controls (*n* = 4, static and dynamic). Values are shown as mean \pm standard error. OcS/BS (*P* < 0.005), NOc/BPm (*P* < 0.005), and BFR/TV (*P* < 0.05) were significantly lower in the *BLG-Cre/PTHrP^{lox/-}* mice than in controls. OcS/BS, osteoclast surface as a percentage of bone surface; NOc/BPm, number of osteoclasts per mm bone perimeter; ObS/BS, osteoblast surface as a percentage of bone surface; NOb/BPm, number of osteoblasts per mm bone perimeter; BFR/TV, bone formation rate as a function of tissue volume; BV/TV, trabecular bone volume as a function of tissue volume; TbTh, trabecular thickness; TbSp, trabecular separation; TbN, trabecular number.